
Introduction to Manifolds

Assignment 2

Due Date: 30/08/2018

Problem 1: In each of the following questions show that the given function f is one-to-one on the given set A . Sketch A and $B = f(A)$. For $p \in B$ find $Df^{-1}(p)$.

1. $f(x, y) = (x^2 - y^2, 2xy)$, $A = \{(x, y) \mid x > 0\}$ and $p = (0, 1)$.
2. $f(x, y) = (e^x \cos y, e^x \sin y)$, $A = \{(x, y) \mid 0 < y < 2\pi\}$ and $p = (0, 1)$.

Problem 2: Let $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be given by the equation $f(x) = \|x\|^2 \cdot x$.

1. Show that f is a smooth function.
2. Show that f maps the unit ball onto itself in a one-to-one fashion.
3. Is the inverse function differentiable at the origin? Why?

Problem 3: Let $g : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be given by $g(x, y) = (2ye^{2x}, xe^y)$ and $f : \mathbb{R}^2 \rightarrow \mathbb{R}^3$ be given by the equation $f(x, y) = (3x - y^2, 2x + y, xy + y^3)$.

1. Show that there is a neighborhood of $(0, 1)$ that g carries in a one-to-one fashion onto a neighborhood of $(2, 0)$.
2. Find $D(f \circ g^{-1})$ at $(2, 0)$.

Problem 4: Let $U \subset \mathbb{R}^n$ be open; let $f : U \rightarrow \mathbb{R}^n$ be a smooth function; assume $Df(x)$ is non-singular for $x \in U$. Show that even if f is not one-to-one on U the image is open in \mathbb{R}^n .

Problem 5: Compute the derivative of the following functions.

1. $f : M(n, \mathbb{R}) \times M(n, \mathbb{R}) \rightarrow M(n, \mathbb{R})$ given by $f(A, B) = A + B$.
2. $g : M(n, \mathbb{R}) \times M(n, \mathbb{R}) \rightarrow M(n, \mathbb{R})$ given by $g(A, B) = AB$.
3. $h : M(n, \mathbb{R}) \rightarrow M(n, \mathbb{R})$ given by $h(A) = A^2$.
4. $j : M(n, \mathbb{R}) \rightarrow M(n, \mathbb{R})$ given by $j(A) = A^t$.