

Classical Mechanics 2, Spring 2016 CMI

Problem set 11

Due by the beginning of lecture on Monday April 4, 2016

Force free motion of a rigid body, Euler angles

1. **⟨9⟩** Consider force free motion of a symmetric top with $I_1 = I_2$, as discussed in the lecture. Suppose the axis of the top makes an angle $\theta \neq 0$ with the fixed direction of \mathbf{L} .
 - (a) **⟨6⟩** Find the angle α between the angular velocity vector $\boldsymbol{\Omega}$ and angular momentum vector \mathbf{L} (α is half the opening angle of the cone swept out by $\boldsymbol{\Omega}$). Express α in terms of θ , the principal moments of inertia and the magnitude of angular momentum L . How does α depend on time and L ?
 - (b) **⟨3⟩** Suppose $I_1 \rightarrow I_3$ so that the symmetric top becomes a spherical top. Based on our study of the spherical top, what do you expect to happen to α ? Is this expectation fulfilled by the above formula for α ?
2. **⟨6⟩** Consider force free rotational motion of a symmetric top ($I_1 = I_2 \neq I_3$) described in terms of Euler angles. Let the co-rotating axes $x_1 = x, x_2 = y$, and $x_3 = z$ be chosen along principal axes of inertia. We follow the notation adopted in the lecture and notes. Write the Lagrangian (no need to derive it) and find the momenta conjugate to the Euler angles and identify which are conserved. From the figure, guess which among p_θ, p_ϕ, p_ψ should correspond to the Z component of angular momentum L_Z , and which should correspond to the $x_3 = z$ component L_z .