

Classical Mechanics 2, Spring 2016 CMI

Problem set 2

Due by the beginning of lecture on Monday Jan 18, 2016

Lagrangian

1. **⟨9⟩** Practice with polar coordinates. Consider a particle moving on the x, y plane $z = 0$ in a central potential $V(r)$. The Lagrangian is $L = \frac{1}{2}m(\dot{x}^2 + \dot{y}^2) - V(r)$. Define plane polar coordinates for the particle's location via $x = r \cos \phi, y = r \sin \phi$. Abbreviate $\sin \phi = s, \cos \phi = c$. Recall that the unit vector in the radial direction is $\hat{r} = c\hat{x} + s\hat{y}$ and that linear momentum is $\mathbf{p} = m\dot{x}\hat{x} + m\dot{y}\hat{y}$. The Euler-Lagrange equations in polar coordinates were found to be $m\ddot{r} = m\dot{\phi}^2 - V'(r)$ and $m\dot{r}\ddot{\phi} = -2m\dot{r}\dot{\phi}$.
 - (a) **⟨2⟩** Show that $\dot{\phi} = \frac{1}{r^2}(x\dot{y} - y\dot{x})$.
 - (b) **⟨3⟩** Draw the unit vector in the direction of increasing ϕ , called $\hat{\phi}$, in a diagram. Express $\hat{\phi}$ as a linear combination of \hat{x}, \hat{y} , using the diagram and an appropriate triangle. Choose $0 < \phi < \pi/2$. Check that $\hat{r} \cdot \hat{\phi} = 0$.
 - (c) **⟨2⟩** Suppose we define the angular velocity as $\vec{\omega} = \dot{\phi}\hat{z}$ and a fictitious force $\vec{F}_c = -2\vec{\omega} \times \vec{p}$. Show that the $\hat{\phi}$ component of \vec{F}_c is what appears on the rhs of the Euler-Lagrange equation $m\dot{r}\ddot{\phi} = -2m\dot{r}\dot{\phi}$.
 - (d) **⟨2⟩** \hat{x}, \hat{y} are constant unit vectors, in the sense that they point in the same direction everywhere on configuration space and also at every point along a trajectory: $\frac{d\hat{x}}{dt} = \frac{d\hat{y}}{dt} = 0$. But $\hat{r}, \hat{\phi}$ change direction from place to place. Show that

$$\frac{d\hat{\phi}}{dt} = -\dot{\phi}\hat{r} \quad \text{and} \quad \frac{d\hat{r}}{dt} = \dot{\phi}\hat{\phi} \quad (1)$$

2. **⟨7⟩** Consider a particle whose dynamics is specified by the Lagrangian

$$L(q, \dot{q}) = \frac{1}{2}m\dot{q}^2 + b(q)\dot{q}. \quad (2)$$

Here $b(q)$ is some differentiable function of q .

- (a) **⟨3⟩** Find the momentum conjugate to q and the equation of motion. What sort of motion does the Lagrangian describe?
 - (b) **⟨4⟩** Explain the nature of this particle by examining the principle of extremal action $S = \int_{t_0}^{t_1} L dt$ for this Lagrangian. Can you relate this action to a more familiar one, how do they differ?
3. **⟨4⟩** A particle of charge e moving in 3D space in the presence of a magnetic field is governed by the Lagrangian $L = \frac{1}{2}m\dot{\mathbf{q}}^2 + \frac{e}{c}\mathbf{A} \cdot \dot{\mathbf{q}}$. Here \mathbf{A} is a 'vector potential' with three components A_i and c is the speed of light.
 - (a) **⟨2⟩** Find the momentum conjugate to the position coordinates q_i .
 - (b) **⟨2⟩** What is the physical dimension of the quantity $e\mathbf{A}$?