

SESHADRI CONSTANTS ON SURFACES

KRISHNA HANUMANTHU

1. PRELIMINARIES

By a *surface*, we mean a projective nonsingular variety of dimension 2 over \mathbb{C} . A *curve* C on a surface X is an effective divisor. The group of divisors on X modulo linear equivalence is isomorphic to isomorphism classes of line bundles on X . We use the additive notation of divisors and multiplicative notation of line bundles interchangeably.

We use notation from [1], especially from Chapter 5. A basic reference for much of this material is [2, 1.5, 5.1, 5.2].

Let X be a surface and let $x \in X$. We will often work with the blow up $\pi : X_1 \rightarrow X$ of X at x . We denote the exceptional divisor of π by $E = \pi^{-1}(x)$. Then we have an isomorphism $\text{Div}(X_1) \cong \text{Div}(X) \oplus \mathbb{Z}$, where $\text{Div}(-)$ denotes the group of divisors modulo linear equivalence. The intersection pairing on $\text{Div}(X_1)$ is completely determined by the following (see [1, Proposition 5.3.2]).

- (1) $\pi^*(C) \cdot \pi^*(D) = C \cdot D$ for $C, D \in \text{Div}(X)$;
- (2) $\pi^*(C) \cdot E = 0$ for $C \in \text{Div}(X)$;
- (3) $E^2 = -1$.

The following is a nice argument for (3)¹.

Choose two curves C, D on X which meet transversally at x . Then the strict transforms of C, D are given by $\pi^*(C) - E$ and $\pi^*(D) - E$ respectively (see [1, Proposition 5.3.6]). Then using properties (1) and (2) above, $C \cdot D - 1 = C \cdot D + E^2$.

We recall the Riemann-Roch theorem for surfaces (see [1, Theorem 5.1.6]).

Theorem 1.1 (Riemann-Roch theorem for surfaces). *Let X be a surface and let L be a line bundle on X . Suppose K denotes the canonical line bundle on X . Then $h^0(X, L) - h^1(X, L) + h^2(X, L) = \frac{L \cdot (L - K)}{2} + 1 - h^1(X, \mathcal{O}_X) + h^2(X, \mathcal{O}_X)$.*

We will also use the following theorem frequently (see [1, Theorem 5.1.10]).

Theorem 1.2 (Nakai criterion for ampleness). *A line bundle L on X is ample if and only if $L^2 > 0$ and $L \cdot C > 0$ for all curves C on X .*

Theorem 1.3 (Kleiman's theorem). *Let X be a surface. Let D be a divisor on X such that $D \cdot C \geq 0$ for every curve on X . Then $D^2 \geq 0$.*

Date: February 4, 2017.

¹We learned this from Pramathanath Sastry.

Proof. Fix an ample divisor H on X . For $t \in \mathbb{R}$, consider the quadratic polynomial $P(t) = (D + tH)^2 = D^2 + 2tD \cdot H + t^2H^2$. We are done if $P(0) \geq 0$. Suppose that $P(0) < 0$.

Since H is ample, there is a curve C in the linear system mH for some $m > 0$. By hypothesis on D , $D \cdot C = D \cdot mH \geq 0$, and hence $D \cdot H \geq 0$. Since the coefficient of t in $P(t)$ is non-negative and the coefficient of t^2 is positive, there is a real number $t_0 > 0$ such that $P(t_0) = 0$.

If $t > t_0$ is a rational number then $D + tH$ is ample. Indeed, this divisor meets all curves positively. Further $P(t) = (D + tH)^2 > 0$, so $D + tH$ is ample by Nakai criterion Theorem 1.2.

Now define $Q(t) = D \cdot (D + tH)$ and $R(t) = tH \cdot (D + tH)$, so that $P(t) = Q(t) + R(t)$ for every $t \in \mathbb{R}$. If $t > t_0$ is a rational number, $Q(t) \geq 0$ since $D + tH$ is ample. Thus $Q(t_0) \geq 0$ too, by continuity. On the other hand, $R(t_0) > 0$. We thus get $P(t_0) > 0$, which is a contradiction. \square

Definition 1.4. A line bundle L is called *nef* if $L \cdot C \geq 0$ for every curve C on X .

2. SESHADRI CONSTANTS

Definition 2.1 (Seshadri constants). Let X be a surface and let L be a nef line bundle on X . The *Seshadri constant of L at a point $x \in X$* , denoted $\epsilon(X, L, x)$, is defined as:

$$\epsilon(X, L, x) := \inf_{x \in C} \frac{L \cdot C}{\text{mult}_x C}.$$

Remark 2.2. It suffices to consider only integral curves in Definition 2.1. Indeed, it is easy to check that for two curves C and D passing through x , we have

$$\frac{L \cdot (C + D)}{\text{mult}_x(C) + \text{mult}_x(D)} \geq \min \left(\frac{L \cdot C}{\text{mult}_x(C)}, \frac{L \cdot D}{\text{mult}_x(D)} \right).$$

Lemma 2.3. Let $\pi : X_1 \rightarrow X$ be the blow up of X at x with exceptional divisor E . Then

$$\epsilon(X, L, x) = \sup \{ \lambda \in \mathbb{R} \mid \pi^*(L) - \lambda E \text{ is nef} \}.$$

Proof. Let ϵ_1 denote the supremum in the lemma and let $\epsilon = \epsilon(X, L, x)$.

Suppose that $\pi^*(L) - \lambda E$ is nef for some λ and let C be a curve with multiplicity m at x . Then the strict transform of C is $\pi^*(C) - mE$. Hence $(\pi^*(L) - \lambda E) \cdot (\pi^*(C) - mE) = L \cdot C - \lambda m \geq 0$. So $\lambda \leq \frac{L \cdot C}{m}$. This shows that $\epsilon_1 \leq \epsilon$. On the other hand, we claim that $\pi^*(L) - \epsilon E$ is nef which proves that $\epsilon \leq \epsilon_1$ and the lemma is proved.

Let C_1 be an integral curve on X_1 . If $C_1 = E$ then $(\pi^*(L) - \epsilon E) \cdot E = \epsilon > 0$. Otherwise, C_1 is the strict transform of $C = \pi(C_1)$. So $C_1 = \pi^*(C) - mE$, where m is the multiplicity of C at x . By definition, $\epsilon \leq \frac{L \cdot C}{m}$ which implies that $(\pi^*(L) - \epsilon E) \cdot C_1 \geq 0$. \square

Theorem 2.4 (Seshadri criterion for ampleness). *A nef line bundle L on X is ample if and only if $\epsilon(X, L, x) > 0$ for every $x \in X$.*

Proof. Let D be a divisor on X corresponding to the line bundle L . If D is ample then a multiple mD is very ample for some positive integer m . Then mD is a hyperplane section for the embedding given by mD . If C is a curve on X passing through x , $mD \cdot C \geq \text{mult}_x(C)$ by properties of intersection numbers. Here we choose a hyperplane section which meets C properly. Hence $\epsilon = \epsilon(X, L, x) \geq \frac{1}{m}$.

Now assume that $\epsilon > 0$. We use Nakai criterion to show that D is ample. The hypothesis clearly implies that $D \cdot C > 0$ for any curve on X . We have to show that $D^2 > 0$. Choose an integer $a > 1/\epsilon$. We claim that $a\pi^*(D) - E$ is nef on the blow up X_1 of X at x . Then by Theorem 1.3, $(a\pi^*(D) - E)^2 = a^2D^2 - 1 \geq 0$, which proves that $D^2 > 0$.

In order to show that $a\pi^*(D) - E$ is nef, we show that it meets all integral curves non-negatively. Since its intersection number with E is 1, we consider a curve $C_1 \neq E$ on X_1 . Then C_1 is the strict transform of $C = \pi(C_1)$. So we may write $C_1 = \pi^*(C) - mE$ where m is the multiplicity of C at x . So $(a\pi^*(D) - E) \cdot C_1 = aD \cdot C - m \geq 0$, since $aD \cdot C \geq a\epsilon m \geq m$. \square

Using one of the directions of the proof of Theorem 2.4, we obtain the following.

Corollary 2.5. *If L is very ample then $\epsilon(X, L, x) \geq 1$ for any $x \in X$.*

Remark 2.6. In fact, one can prove that $\epsilon(X, L, x) \geq 1$ for any $x \in X$ if L is merely ample and base point free. Indeed, let L be such a line bundle. Let $x \in X$ and let C be an irreducible and reduced curve on X passing through x . Then there exists a curve D in the linear system $|L|$ such that $x \in D$ and C is not a component of D . This is because the sections of L determine a morphism $f : X \rightarrow \mathbb{P}^n$ and we may take a hyperplane $H \subset \mathbb{P}^n$ containing $f(x)$. Since L is ample, the morphism f is finite. In particular, $f(C)$ is not a point. So the pull-back D of H has the required properties. Now $L \cdot C = D \cdot C \geq \text{mult}_x(C)$.

Proposition 2.7 (An upper bound). *Given X, L, x as in Definition 2.1, we have $\epsilon(X, L, x) \leq \sqrt{L^2}$.*

Proof. We use the equivalent formulation of $\epsilon = \epsilon(X, L, x)$ in Lemma 2.3. With the notation in Lemma 2.3, $\pi^*(L) - \epsilon E$ is nef. By Theorem 1.3, $0 \leq (\pi^*(L) - \epsilon E)^2 = L^2 - \epsilon^2$. \square

Example 2.8. Let $p, q \in \mathbb{P}^2$ and let $\pi : X \rightarrow \mathbb{P}^2$ be the blow up of \mathbb{P}^2 at p, q . Let H be the pull-back of $\mathcal{O}_{\mathbb{P}^2}(1)$ and let E_1, E_2 be the exceptional curves. Let $L = 3H - E_1 - E_2$. Denote the strict transform of the line through p, q in \mathbb{P}^2 by l .

We claim that $\epsilon(X, L, x) = 1$ if $x \in E_1 \cup E_2 \cup l$ and $\epsilon(X, L, x) = 2$ otherwise.

It is not hard to see that L is ample and base point free (in fact L is very ample). Hence by Remark 2.6, $\epsilon(X, L, x) \geq 1$ for all $x \in X$.

Let $x \in E_1 \cup E_2 \cup l$. Then $\epsilon(X, L, x) = 1$, since $L \cdot E_1 = L \cdot E_2 = L \cdot l = 1$ and E_1, E_2, l are nonsingular.

Let $x \notin E_1 \cup E_2 \cup l$. Let l_1, l_2 be the strict transforms of the lines through $p, \pi(x)$ and $q, \pi(x)$. Since $L \cdot l_1 = 2$, we have $\epsilon(X, L, x) \leq 2$. We now show that $\epsilon(X, L, x) \geq 2$, to complete the argument.

Let C be any irreducible and reduced curve on X passing through x . Assume that C is different from E_1, E_2, l, l_1, l_2 . Then C is the strict transform of $\pi(C)$ and we may write $C = dH - m_1E_1 - m_2E_2$, where d is the degree of $\pi(C)$ and m_i are the multiplicities of $\pi(C)$ at p, q . Let m be the multiplicity of $\pi(C)$ at $\pi(x)$. Since $d > 1$ by assumption, we have $C \cdot l \geq 0, C \cdot l_1 \geq 0$ and $C \cdot l_2 \geq 0$. Hence $d \geq m_1 + m_2, d \geq m + m_1$ and $d \geq m + m_2$. This gives $3d \geq 2m + m_1 + m_2$, which in turn implies that $L \cdot C \geq 2m$. Hence $\epsilon(X, L, x) \geq 2$.

Definition 2.9. Let X be a surface.

- (1) Let $x \in X$. The *Seshadri constant of X at x* is defined as $\epsilon(X, x) = \inf \epsilon(X, L, x)$, where the infimum is taken over all ample line bundles L on X .
- (2) Let L be an ample line bundle on X . The *Seshadri constant of L* is defined as $\epsilon(X, L) = \inf \epsilon(X, L, x)$, where the infimum is taken over all points $x \in X$.
- (3) The *Seshadri constant of X* is defined as $\epsilon(X) = \inf \epsilon(X, L)$, where the infimum is taken over all ample line bundles L on X . Equivalently, $\epsilon(X) = \inf \epsilon(X, x)$, where the infimum is taken over all points $x \in X$.

Question 2.10 (Some open questions about Seshadri constants).

- (1) Is there a surface X such that $\epsilon(X) = 0$?
- (2) Is there a triple (X, L, x) such that $\epsilon(X, L, x)$ is irrational?

However the Seshadri constants can be arbitrarily small when we vary the surface.

Theorem 2.11 (Miranda's example). *Given a real number $\delta > 0$ there exists a surface X and an ample line bundle L on X such that $\epsilon(X, L, x) < \delta$ for some $x \in X$.*

Proof. Choose $m \in \mathbb{N}$ so that $m > \frac{1}{8}$. For some $d > 2$, let $\Gamma \in \mathbb{P}^2$ be an integral curve of degree d with a point $x \in \mathbb{P}^2$ of multiplicity m . Now choose another integral curve Γ' meeting Γ transversally such that the linear system spanned by Γ and Γ' consists of only integral curves. Such a Γ' exists by Lemma 2.12 below.

Let $\pi : X \rightarrow \mathbb{P}^2$ be the blow up of \mathbb{P}^2 at $\Gamma \cap \Gamma' = \{p_1, \dots, p_{d^2}\}$. Denote by C and C' the strict transforms of Γ and Γ' respectively. Since p_i are all simple points on Γ and Γ' , C, C' are isomorphic to Γ, Γ' , respectively. In particular, C contains a point $x \in X$ of multiplicity m . Moreover the pencil $\langle C, C' \rangle$ spanned by C and C' consists of only integral curves.

We have a morphism $f : X \rightarrow \mathbb{P}^1$ determined by the linear system $\langle C, C' \rangle$. Note that f resolves the indeterminacies of the map $\mathbb{P}^2 \dashrightarrow \mathbb{P}^1$ determined by the pencil $\langle \Gamma, \Gamma' \rangle$. Let E be an exceptional divisor on X for the blow up π . For some $a \geq 2$, let $L = aC + E$. We claim that L is ample.

Since $C^2 = 0, C \cdot E = 1$ and $E^2 = -1$, we have $L^2 = 2a - 1 > 0, L \cdot C = 1 > 0$ and $L \cdot E = a - 1 > 0$. By Nakai, it suffices to show that $L \cdot D > 0$ for every curve D on X .

If D is a fibre of f then D is inside a curve in the pencil $\langle C, C' \rangle$. But since all curves in this linear system are integral, D itself is in it. So D is linearly equivalent to C and $L \cdot D = 1$. If D is not a fibre of f , let $p \in \mathbb{P}^1$. Since $f^{-1}(p)$ is a curve in the pencil $\langle C, C' \rangle$, it is linearly equivalent to C . As D is not in this linear system and it meets $f^{-1}(p)$ in a nonempty set $f^{-1}(p) \cap D$, we have $C \cdot D > 0$. Further it is clear that $D \cdot E \geq 0$. Indeed, if $D \cdot E < 0$ then E

is a component of D . Since D is integral it follows that $D = E$. Thus $L \cdot D \geq \alpha - 1 > 0$ and L is ample.

Finally, $\epsilon(X, L, x) \leq \frac{L \cdot C}{\text{mult}_x(C)} = \frac{1}{m} < \delta$. \square

Lemma 2.12. *Let $\Gamma \subset \mathbb{P}^2$ be an integral curve of degree $d > 2$. Then there exists an integral curve Γ' which intersects Γ transversally such that the linear system of plane curves spanned by Γ and Γ' contains only integral curves.*

Proof. The (projective) space of degree d plane curves has dimension $\binom{d+2}{2} - 1$. A degree d curve C is not integral precisely when there are polynomials f_1, f_2 such that $f = f_1 f_2$ where f is the form defining C . Thus the space of all non-integral degree d curves has dimension bounded above by $\binom{e+2}{2} + \binom{d-e+2}{2} - 2 = \frac{d^2+3d}{2} + e^2 - de$, for $e \in \{1, 2, \dots, d-1\}$. This number is largest when $e = 1$ and in this case it is equal to $\frac{d^2+3d}{2} + 1 - d$. So the space of all non-integral degree d curves has codimension at least 1 in the space of all degree d curves. So a ‘‘general line’’ through Γ avoids this subspace. More precisely, there is a dense open set of integral curves Γ' such that the pencil spanned by Γ and Γ' consists only of integral curves.

Now by Bertini’s theorem, there is a dense open subspace of degree d curves consisting of integral curves which meet Γ transversally. So we may choose Γ' satisfying the desired properties. \square

In view of Proposition 2.7, we say that a Seshadri constant $\epsilon(X, L, x)$ is *sub-optimal* if $\epsilon(X, L, x) < \sqrt{L^2}$. We further say that a curve C is a *Seshadri curve* if $\epsilon(X, L, x) = \frac{L \cdot C}{\text{mult}_x(C)}$. The following theorem says that there is always a Seshadri curve for a sub-optimal Seshadri constant. As a consequence, they are always rational.

Theorem 2.13. *Let X be a surface and let L be an ample line bundle on X . Let $x \in X$. Suppose that $\epsilon(X, L, x) < \sqrt{L^2}$. Then there is an irreducible curve C on X such that $\epsilon(X, L, x) = \frac{L \cdot C}{\text{mult}_x(C)}$.*

Proof. Denote $\epsilon = \epsilon(X, L, x)$. Let $\mu \in \mathbb{R}$ be such that $1 < \mu < \frac{\sqrt{L^2}}{\epsilon}$.

By definition of ϵ , there is a sequence $\{C_n\}_{n \geq 1}$ of irreducible curves such that $\frac{L \cdot C_n}{\text{mult}_x(C_n)}$ converges to ϵ . Since $\epsilon < \frac{\sqrt{L^2}}{\mu}$, there exists some N such that $\frac{L \cdot C_n}{\text{mult}_x(C_n)} < \frac{\sqrt{L^2}}{\mu}$ for $n \geq N$.

Next we claim that for some $d > 0$, there exists a divisor $D \in |dL|$ such that $\frac{L \cdot D}{\text{mult}_x(D)} \leq \mu \sqrt{L^2}$. By Riemann-Roch Theorem 1.1, $h^0(dL) \geq \frac{d^2 L^2}{2}$ for large d . Since a point of multiplicity m imposes $\binom{m+1}{2}$ conditions, there is a curve D in the linear system $|dL|$ with a point of multiplicity m if $d^2 L^2 \geq m^2 + m$. Clearly this holds if $d \geq \frac{m}{\sqrt{L^2}} + 1$. Hence, by taking large enough m , we can make the ratio $\frac{d}{m}$ arbitrarily close to $\frac{1}{\sqrt{L^2}}$. In other words, the ratio $\frac{L \cdot D}{\text{mult}_x(D)} = \frac{dL^2}{\text{mult}_x(D)}$ is arbitrarily close to $\sqrt{L^2}$. Hence we conclude that for some $d > 0$, there exists a divisor $D \in |dL|$ such that $\frac{L \cdot D}{\text{mult}_x(D)} \leq \mu \sqrt{L^2}$, since $\mu > 1$.

The theorem now follows by the lemma below. The lemma implies that the collection of curves $\{C_n\}$ is finite. So the limit ϵ is achieved by some C_n . \square

Lemma 2.14. *Given a real number $\mu > 0$, suppose that there is a divisor $D \in |dL|$ for some $d > 0$, such that $\frac{L \cdot D}{\text{mult}_x(D)} \leq \mu \sqrt{L^2}$. Then every irreducible curve C satisfying $\frac{L \cdot C}{\text{mult}_x(C)} < \frac{\sqrt{L^2}}{\mu}$ is a component of D .*

Proof. If C is not a component of D , then C and D meet properly. By Bezout's theorem, $D \cdot C = dL \cdot C \geq (\text{mult}_x(D))(\text{mult}_x(C)) > \frac{L \cdot D}{\mu \sqrt{L^2}} \frac{\mu L \cdot C}{\sqrt{L^2}} = dL \cdot C$. This is a contradiction. \square

2.1. Multi-point Seshadri constants. Let X be a smooth projective surface and let L be a nef line bundle on X . Let $r \geq 1$ be an integer. For $x_1, \dots, x_r \in X$, the *multi-point Seshadri constant* of L at x_1, \dots, x_r is defined as follows.

$$\epsilon(X, L, x_1, \dots, x_r) := \inf_{C \cap \{x_1, \dots, x_r\} \neq \emptyset} \frac{L \cdot C}{\sum_i^r \text{mult}_{x_i} C}.$$

As in case of the single point Seshadri constants, we note that the above infimum is the same as the infimum taken over irreducible, reduced curves C such that $C \cap \{x_1, \dots, x_r\} \neq \emptyset$.

We have the following proposition which is an analogue of Proposition 2.7.

Proposition 2.15 (An upper bound). *Given X, L, x_1, \dots, x_r as above, we have $\epsilon(X, L, x_1, \dots, x_r) \leq \sqrt{\frac{L^2}{r}}$.*

Definition 2.16. Let X, L, r be as above. We set $\epsilon(X, L, r) := \max_{x_1, \dots, x_r \in X} \epsilon(X, L, x_1, \dots, x_r)$.

The value $\epsilon(X, L, r)$ is attained at a *very general* set of points $x_1, \dots, x_r \in X$. For a reference, see [4]. Here *very general* means that (x_1, \dots, x_r) is outside a countable union of proper Zariski closed sets in $X^r = X \times X \times \dots \times X$.

3. EXAMPLES

3.1. $(X, L) = (\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(1))$. We first consider the case of projective plane in detail.

If $r = 1$, it is easy to see that $\epsilon(X, L, x) = 1$ for any $x \in X$. Similarly, if $r = 2$ and x_1, x_2 are arbitrary points on X , then $\epsilon(X, L, x_1, x_2) = 1/2$.

Now let $x_1, x_2, x_3 \in X$. If the points are collinear, then the line through them is a Seshadri curve and $\epsilon(X, L, x_1, x_2, x_3) = 1/3$. On the other hand, if the points are not collinear, one has $\epsilon(X, L, x_1, x_2, x_3) = 1/2$. Indeed, let l be the line through x_1 and x_2 . By hypothesis, $x_3 \notin l$. Hence the Seshadri quotient for l is $1/2$. Let $C \neq l$ be any other irreducible curve of degree d passing through at least one of the points x_1, x_2 or x_3 . For $i = 1, 2, 3$, let m_i denote the multiplicity of C at x_i . It is clear that $d \geq m_i$ for every i . Applying Bezout to the irreducible curves C and l , we have $d \geq m_1 + m_2$. Hence we obtain $2d \geq m_1 + m_2 + m_3$. Hence $\epsilon(X, L, x_1, x_2, x_3) = 1/2$.

Thus we have $\epsilon(X, L, 3) = 1/2$, and this is achieved for any three non-collinear points.

Similarly, it is not hard to prove that $\epsilon(X, L, 4) = 1/2$, $\epsilon(X, L, 5) = \epsilon(X, L, 6) = 2/5$, $\epsilon(X, L, 7) = 3/8$, $\epsilon(X, L, 8) = 6/17$ and $\epsilon(X, L, 9) = 3$.

For $r \geq 10$, the value of $\epsilon(X, L, r)$ is not known except when r is a square. If $r = s^2$ and $s \geq 3$, Nagata [3] showed that $\epsilon(X, L, r) = 1/s$. In the same paper, Nagata made the following famous conjecture.

Conjecture 3.1 (Nagata Conjecture). $\epsilon(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(1), r) = \frac{1}{\sqrt{r}}$, when $r \geq 10$.

Note that an affirmative solution to Nagata Conjecture for even one non-square value of $r \geq 10$ also solves Question 2.10 (2).

Nagata Conjecture 3.1 is a statement about linear systems of plane curves in the sense we explain now.

Let x_1, \dots, x_r be very general points of \mathbb{P}^2 . Let d, m_1, \dots, m_r be non-negative integers. We are interested in the question:

Is there a curve $C \in \mathbb{P}^2$ of degree d passing through x_i with multiplicity at least m_i for each $1 \leq i \leq r$?

Let $\mathcal{L}(d, m_1, \dots, m_r)$ denote the linear system of plane curves of degree d passing through x_i with multiplicity at least m_i for each $1 \leq i \leq r$. So our question can be rephrased as:

Is $\mathcal{L}(d, m_1, \dots, m_r)$ non-empty?

The linear system of degree d plane curves has dimension $\frac{d(d+3)}{2}$. This can be seen by observing that a degree d plane curve is given by a homogeneous form of degree d in three variables. This is also a special case of Riemann-Roch Theorem 1.1.

On the other hand, a point of multiplicity m imposes $\binom{m+1}{2}$ conditions on the linear systems of curves of degree d . It follows that $\mathcal{L}(d, m_1, \dots, m_r)$ is non-empty provided $\frac{d(d+3)}{2} \geq \sum_{i=1}^r \binom{m_i+1}{2}$. Let us refer to this as the Riemann-Roch inequality.

It is easy to prove that if d, m_1, \dots, m_r satisfy the Riemann-Roch inequality, then the Seshadri quotient $\frac{d}{\sum_i m_i}$ is at least $\frac{1}{\sqrt{r}}$, as predicted by the Nagata Conjecture. However this inequality is not necessary for existence of such curves, as the example below shows.

Consider $r = 11$, $d = 10$, $m_1 = 8$, and $m_2 = \dots = m_{11} = 2$. Then $\frac{d(d+3)}{2} = 65$ while $\sum_{i=1}^{11} \binom{m_i+1}{2} = 66$. However there is a curve in the linear system $\mathcal{L}(10, 8, 2, \dots, 2)$! Indeed, we first observe that $\mathcal{L}(5, 4, 1, \dots, 1)$ is non-empty (again $r = 11$). This follows from Riemann-Roch inequality. Let $C \in \mathcal{L}(5, 4, 1, \dots, 1)$. Then $2C \in \mathcal{L}(10, 8, 2, \dots, 2)$. Note that the Seshadri quotient for $2C$ is $\frac{10}{28} = 0.3571$, while the Nagata Conjecture predicts that $\epsilon(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(1), 11) = \frac{1}{\sqrt{11}} = 0.3015$. The curve C is an example of *exceptional curve*: its proper transform on the blow up of \mathbb{P}^2 at x_1, \dots, x_{11} is a smooth rational curve of self-intersection -1 .

So, in order to prove the Nagata Conjecture, one needs to understand effective curves given by d, m_1, \dots, m_r that do *not* satisfy the Riemann-Roch inequality.

There is a precise conjecture, called the *SHGH Conjecture*, which completely characterizes tuples (d, m_1, \dots, m_r) which represent effective plane curves. It says, in part, that any effective linear system of plane curves given by a tuple (d, m_1, \dots, m_r) not satisfying the Riemann-Roch inequality contains a non-reduced multiple of an exceptional curve. The SHGH Conjectures implies the Nagata Conjecture.

REFERENCES

- [1] Hartshorne, Robin, *Algebraic Geometry*.
- [2] Lazarsfeld, Robert, *Positivity in Algebraic Geometry I*.
- [3] Nagata, Masayoshi, *On the 14-th problem of Hilbert*, Am. J. Math. 81 (1959), 766-772.
- [4] Oguiso, Keiji, *Seshadri constants in a family of surfaces*, Math. Ann. 323 (2002), no. 4, 625-631.