

DOUBLE COMPLEXES

We work throughout in an abelian category \mathcal{A} in which *countable direct sums* exist.

There are two related notions of a double complex. We give both versions below. The second (i.e., what we call an *anti-commuting* double complex below) is what you will often find in the older literature—and amongst non-algebraic-geometers. The first version (which we simply call a double complex, or sometimes a *standard double complex*) is the version given by Grothendieck in EGA, and is what most algebraic geometers are used to. The difference is one of convention.

Standard Double Complexes. A *double complex* in \mathcal{A} , or sometimes in our class, a *standard double complex* in \mathcal{A} , consists of data $A^{\bullet\bullet} = (A, \partial_1, \partial_2)$, where

$$A = (A^{p,q})_{(p,q) \in \mathbb{Z} \times \mathbb{Z}}$$

is a family of objects in \mathcal{A} , and

$$\partial_1 = (\partial_1^{p,q})_{(p,q) \in \mathbb{Z}} \quad \partial_2 = (\partial_2^{p,q})_{(p,q) \in \mathbb{Z}}$$

are two families of morphisms

$$\partial_1^{p,q}: A^{p,q} \rightarrow A^{p+1,q} \quad \partial_2^{p,q}: A^{p,q} \rightarrow A^{p,q+1}$$

such that

$$\partial_1 \partial_1 = 0 \quad \partial_2 \partial_2 = 0 \quad \partial_1 \partial_2 = \partial_2 \partial_1.$$

We often suppress the superscripts p, q when these are either immaterial or easily deducible from the context. Thus, e.g., we write ∂_2 for $\partial_2^{p,q}$. The maps ∂_1 and ∂_2 will be called partial coboundaries, and when we wish to be more specific, they will be called *horizontal* and *vertical* (partial) coboundaries respectively. The data fits into a commutative diagram, whose rows and columns are complexes.

$$\begin{array}{ccccccc}
 & & \vdots & & \vdots & & \vdots \\
 & & \uparrow \partial_2 & & \uparrow \partial_2 & & \uparrow \partial_2 \\
 \dots & \xrightarrow{\partial_1} & A^{0,q+1} & \xrightarrow{\partial_1} & \dots & \xrightarrow{\partial_1} & A^{p,q+1} & \xrightarrow{\partial_1} & A^{p+1,q+1} & \xrightarrow{\partial_1} & \dots \\
 & & \uparrow \partial_2 & & \uparrow \partial_2 & & \uparrow \partial_2 & & \uparrow \partial_2 \\
 \dots & \xrightarrow{\partial_1} & A^{0,q} & \xrightarrow{\partial_1} & \dots & \xrightarrow{\partial_1} & A^{p,q} & \xrightarrow{\partial_1} & A^{p+1,q} & \xrightarrow{\partial_1} & \dots \\
 & & \uparrow \partial_2 & & \uparrow \partial_2 & & \uparrow \partial_2 & & \uparrow \partial_2 \\
 & & \vdots & & \vdots & & \vdots & & \vdots
 \end{array}$$

Next consider the direct sum¹

$$\mathrm{Tot}^n A^{\bullet\bullet} := \bigoplus_{p+q=n} A^{p,q}.$$

¹This is where our assumption that \mathcal{A} has countable direct sums comes into play. Alternately, one can assume that the displayed direct sum for $\mathrm{Tot}^n A^{\bullet\bullet}$ is finite for every $n \in \mathbb{Z}$.

Define

$$\partial^n: \text{Tot}^n A^{\bullet\bullet} \rightarrow \text{Tot}^{n+1} A^{\bullet\bullet}$$

by the formula

$$\partial^n = \sum_{p+q=n} \{\partial_1^{p,q} + (-1)^p \partial_2^{p,q}\}.$$

The map within “curly brackets” can be regarded as a map $A^{p,q} \rightarrow \text{Tot}^{n+1} A^{\bullet\bullet}$, taking values in the subobject $A^{p+1,q} \oplus A^{p,q+1}$ of $\text{Tot}^{n+1} A^{\bullet\bullet}$, whence by the definition of direct sum, the map ∂^n makes sense.

Evidently

$$\partial^{n+1} \circ \partial^n = 0$$

for every $n \in \mathbb{Z}$ by the relations given between ∂_1 and ∂_2 . We have therefore a complex $(\text{Tot}^\bullet A^{\bullet\bullet}, \partial)$, called the *total complex* associated to the double complex $A^{\bullet\bullet}$.

A morphism of double complexes $\varphi: A^{\bullet\bullet} \rightarrow B^{\bullet\bullet}$ is (of course) a family of maps $f^{p,q}: A^{p,q} \rightarrow B^{p,q}$, one for each ordered pair of integers (p, q) , which commute with vertical and horizontal coboundaries. This naturally induces a map of complexes $\text{Tot} f: \text{Tot}^\bullet A^{\bullet\bullet} \rightarrow \text{Tot}^\bullet B^{\bullet\bullet}$

Anti-commutative double complexes. In much of the pre-Grothendieck literature, double complexes mean a variant of our standard double complexes. The only difference is that the grids in the diagram on the last page anti-commute rather than commute. In greater detail, for this course, data of the form $K^{\bullet\bullet} = (K, d_1, d_2)$ represents an *anti-commuting* double complex if K is a family $(K^{p,q})$ of objects in \mathcal{A} indexed by $\mathbb{Z} \times \mathbb{Z}$ and $d_1 = (d_1^{p,q}: K^{p,q} \rightarrow K^{p+1,q})$, $d_2 = (d_2^{p,q}: K^{p,q} \rightarrow K^{p,q+1})$ are families of maps indexed by $(p, q) \in \mathbb{Z} \times \mathbb{Z}$, called the *horizontal* and *vertical* partial coboundaries respectively, such that

$$d_1 d_1 = 0 \quad d_2 d_2 = 0, \quad d_1 d_2 = -d_2 d_1.$$

We set (and please pay attention to the notation, especially the accent on the top left)

$${}'\text{Tot}^n K^{\bullet\bullet} := \bigoplus_{p+q=n} K^{p,q}$$

and define

$$d^n: {}'\text{Tot}^n K^{\bullet\bullet} \rightarrow {}'\text{Tot}^{n+1} K^{\bullet\bullet}$$

by the formula

$$d^n = \sum_{p+q=n} (d_1^{p,q} + d_2^{p,q})$$

without any sign of the form $(-1)^p$ intervening. It is easy to see, with $d := (d^n)_{n \in \mathbb{Z}}$, that $({}'\text{Tot}^\bullet K^{\bullet\bullet}, d)$ is a complex. We call this complex the total complex associated with the anti-commuting double complex $K^{\bullet\bullet}$.

I will leave the task of defining maps of anti-commuting double complexes to you.

Bounded double complexes. Let $C^{\bullet\bullet}$ be a double complex (standard or anti-commutative). We say it is *bounded on the left* if there is an integer p_0 such that

$$C^{p,q} = 0, \quad p < p_0.$$

If this happens we sometimes say $C^{\bullet\bullet}$ is *bounded on the left by p_0* . Similarly $C^{\bullet\bullet}$ is *bounded below (by q_0)* if there exists an integer q_0 such that

$$C^{p,q} = 0 \quad q < q_0.$$

I leave to you the fun task of defining terms like *bounded on the right* and *bounded above*.

Note that if $C^{\bullet\bullet}$ is bounded on the left and below (resp. above and to the right) it lives in a translate of the first quadrant (resp. third quadrant) and as such the direct sum

$$\bigoplus_{p+q=n} C^{p,q}$$

is actually a finite sum² for each n . So in such instances, one can define $\text{Tot}^n C^{\bullet\bullet}$ or ${}^{\prime}\text{Tot}^n C^{\bullet\bullet}$ (as the case may be) without insisting that \mathcal{A} have countable direct sums. In fact we will largely be dealing with such situations.

²Draw a picture with such quadrant translates, and look at the intersection of such quadrant translates with lines having slope -1 .