

**Definition 1 (Neighbourhood equivalence)** Two valuations  $v$  and  $v'$  are said to be neighbourhood equivalent, written as  $v \simeq_{\text{nbd}} v'$  if:

1.  $\lfloor v(x) \rfloor = \lfloor v'(x) \rfloor$  for all clocks  $x$ ,
2.  $\{v(x)\} = 0$  iff  $\{v'(x)\} = 0$  for all clocks  $x$
3. for every pair of clocks  $x, y$ :
  - (a)  $\{v(x)\} < \{v(y)\} \Leftrightarrow \{v'(x)\} < \{v'(y)\}$
  - (b)  $\{v(x)\} = \{v(y)\} \Leftrightarrow \{v'(x)\} = \{v'(y)\}$

Each equivalence class of  $\simeq_{\text{nbd}}$  will be called a *neighbourhood*.

---

1. Group the following valuations over five clocks into neighbourhoods.

$$\begin{aligned}
 v_1 &:= (7.4, 2.1, 8.7, 5.4, 7.0) \\
 v_2 &:= (3.4, 2.0, 8.5, 10.0, 7.1) \\
 v_3 &:= (7.3, 2.2, 8.8, 5.2, 7.0) \\
 v_4 &:= (7.5, 2.1, 8.9, 5.5, 7.0) \\
 v_5 &:= (3.2, 2.0, 8.8, 10.0, 7.5) \\
 v_6 &:= (3.3, 2.0, 8.4, 10.0, 7.2)
 \end{aligned}$$

2. Below are combinations of  $v, \delta$  and  $v'$  with  $v \simeq_{\text{nbd}} v'$ . For each of them, find a  $\delta' \geq 0$  such that  $v + \delta \simeq_{\text{nbd}} v' + \delta'$ .

$$\begin{array}{lll}
 v = (1.1, 2.0, 3.0, 4.0, 5.9) & \delta = 3.4 & v' = (1.5, 2.0, 3.0, 4.0, 5.6) \\
 v = (10.2, 4.8, 19.1, 2.0, 8.5) & \delta = 5.4 & v' = (10.1, 4.9, 19.05, 2.0, 8.7)
 \end{array}$$

Recall that  $v + \delta$  is the valuation obtained by adding  $\delta$  to each coordinate of  $v$ .